




News - News Topic: Physics



FASER measures high-energy neutrino interaction strength

The interaction strength of neutrinos had never previously been measured in this energy range

4 APRIL, 2024 | By Piotr Traczyk ([/authors/piotr-traczyk](https://home.cern/authors/piotr-traczyk))



The FASER detector installed 480 m away from the ATLAS detector collision point, in the LHC side tunnel TI12. (image: Maximilien Brice/CERN)

Operating at CERN’s Large Hadron Collider (LHC) since 2022, the [FASER](https://home.cern/science/experiments/faser) (<https://home.cern/science/experiments/faser>) experiment is designed to search for extremely weakly interacting particles. Such particles are predicted by many theories beyond the Standard Model that are attempting to solve outstanding problems in physics such as the nature of dark matter and the matter-antimatter imbalance in the Universe. Another goal of the experiment is to study interactions of high-energy neutrinos produced in the LHC collisions, particles that are nearly impossible to detect in the four big LHC experiments. Last week, at the annual (<http://moriond.in2p3.fr/>) [Rencontres de Moriond](http://moriond.in2p3.fr/) (<http://moriond.in2p3.fr/>) conference, the FASER collaboration presented a measurement of the interaction strength, or “cross section”, of electron neutrinos (ν_e) and muon neutrinos (ν_μ). This is the first time such a measurement has been made at a particle collider. Measurements of this kind can provide important insights across different aspects of physics, from understanding the production of “forward” particles in the LHC collisions and improving our understanding of the structure of the proton to interpreting measurements of high-energy neutrinos from astrophysical sources performed by neutrino-telescope experiments.

Related Articles

[\(/news/news/physics/shine-shines-light-neutrino-beams\)](https://home.cern/news/news/physics/shine-shines-light-neutrino-beams)

[\(/news/news/physics/shine-shines-light-neutrino-beams\)](https://home.cern/news/news/physics/shine-shines-light-neutrino-beams)

Physics | News | 7 June, 2023

[\(/news/news/experiments/live-particle-pursuit-journey-deep-underground-neutrino-experiments\)](https://home.cern/news/news/experiments/live-particle-pursuit-journey-deep-underground-neutrino-experiments)

[\(/news/news/experiments/live-particle-pursuit-journey-deep-underground-neutrino-experiments\)](https://home.cern/news/news/experiments/live-particle-pursuit-journey-deep-underground-neutrino-experiments)

Experiments | News | 6 June, 2023

[\(/news/news/physics/cms-tries-out-seesaw\)](https://home.cern/news/news/physics/cms-tries-out-seesaw)

[\(/news/news/physics/cms-tries-out-seesaw\)](https://home.cern/news/news/physics/cms-tries-out-seesaw)


Physics | News | 4 May, 2023

[View all news >](#)

FASER is located in a side tunnel of the LHC accelerator, 480 metres away from the ATLAS detector collision point. At that location, the LHC beam is already nearly 10 metres away, bending away on its circular 27-kilometre path. This is a unique location for studying weakly interacting particles produced in the LHC collisions. Charged particles produced in the collisions are deflected by the LHC magnets. Most neutral particles are stopped by the hundreds of metres of rock between FASER and ATLAS. Only very weakly interacting neutral particles like neutrinos are expected to continue straight on and reach the location where the detector is installed.

The probability of a neutrino interacting with matter is very small, but not zero. The type of interaction that FASER is sensitive to is where a neutrino interacts with a proton or a neutron inside the detector. In this interaction, the neutrino transforms into a charged “lepton” of the same family – an electron in the case of an ν_e , and a muon in the case of a ν_μ – which is visible in the detector. If the energy of the neutrino is high, several other particles are also produced in the collision.

The detector used to perform the measurement consists of 730 interleaved tungsten plates and photographic emulsion plates. The emulsion was exposed during the period from 26 July to 13 September 2022 and then chemically developed and analysed in search of charged particle tracks. Candidates for neutrino interactions were identified by looking for clusters of tracks that could be traced back to a single vertex. One of these tracks then had to be identified as a high-energy electron or muon.



Event displays identified by the FASER collaboration as candidates for an ν_e (left) and a ν_μ (right) interacting in the detector. Invisible here, the neutrinos arrive from the left and then interact to create multiple tracks spraying out to the right (coloured lines), one of which is identified as a charged lepton (labelled). (credit: FASER collaboration)

In total, four candidates for an ν_e interaction and eight candidates for a ν_μ interaction have been found. The four ν_e candidates represent the first direct observation of electron neutrinos produced at a collider. The observations can be interpreted as measurements of neutrino interaction cross sections, yielding $(1.2^{+0.9}_{-0.8}) \times 10^{-38} \text{ cm}^2 \text{ GeV}^{-1}$ in the case of the ν_e and $(0.5 \pm 0.2) \times 10^{-38} \text{ cm}^2 \text{ GeV}^{-1}$ in the case of the ν_μ . The energies of the neutrinos were found to be in a range between 500 and 1700 GeV. No measurement of the neutrino interaction cross section had previously been made at energies above 300 GeV in the case of the ν_e and between 400 GeV and 6 TeV in the case of the ν_μ .

The results obtained by FASER are consistent with expectations and demonstrate the ability of FASER to make neutrino cross-section measurements at the LHC. With the full LHC Run 3 data, 200 times more neutrino events will be detected, allowing much more precise measurements.

Read more in the FASER [publication \(https://arxiv.org/abs/2403.12520\)](https://arxiv.org/abs/2403.12520).

[neutrino \(/tags/neutrino\)](/tags/neutrino) [FASER \(/tags/faser\)](/tags/faser) [LHC experiments \(/tags/lhc-experiments\)](/tags/lhc-experiments)

Also On Physics