

# SuperWIMP Cosmology and Collider Phenomenology


Jonathan Feng  
University of California, Irvine

SUSY04, Tsukuba  
21 June 2004

## Based On...

- Feng, [Rajaraman](#), [Takayama](#), *Superweakly Interacting Massive Particles*, hep-ph/0302215
- Feng, [Rajaraman](#), [Takayama](#), *SuperWIMP Dark Matter Signals from the Early Universe*, hep-ph/0306024
- Feng, [Rajaraman](#), [Takayama](#), *Probing Gravitational Interactions of Elementary Particles*, hep-th/0405248
- Feng, [Su](#), [Takayama](#), *Gravitino Dark Matter from Slepton and Sneutrino Decays*, hep-ph/0404198
- Feng, [Su](#), [Takayama](#), *Supergravity with a Gravitino LSP*, hep-ph/0404231

# Dark Matter



- Tremendous recent progress:  
 $\Omega_{\text{DM}} = 0.23 \pm 0.04$
- But...we have no idea what it is
- Precise, unambiguous evidence for new particle physics

# SuperWIMPs – New DM Candidate

- Why should we care?

We already have axions, warm gravitinos, neutralinos, Kaluza-Klein particles, Q balls, wimpzillas, self-interacting particles, self-annihilating particles, fuzzy dark matter, branons...

- SuperWIMPs have all the virtues of neutralinos...

Well-motivated stable particle

Naturally obtains the correct relic density


Rich implications for cosmology, astrophysics, colliders

- ...and more: There is already a signal

# SuperWIMPs: The Basic Idea


- Supergravity gravitinos: mass  $\sim M_W$ , couplings  $\sim M_W/M_*$

- $\tilde{G}$  not LSP




- Assumption of most of literature


- $\tilde{G}$  LSP



- Completely different cosmology and phenomenology



- Assume  $\tilde{G}$  LSP, WIMP NLSP
- WIMPs freeze out as usual



- But at  $t \sim M_*^2/M_W^3 \sim \text{year}$ , WIMPs decay to gravitinos

Gravitinos are dark matter now: they are *superWIMPs*, superweakly interacting massive particles

# SuperWIMP Virtues

- Well motivated stable particle

Predicted by supersymmetry (with R-parity conservation, high-scale SUSY breaking)

Completely generic – present in “1/2” of parameter space

- Naturally obtains the correct relic density:

$$\Omega_{\tilde{G}} \sim \Omega_{\text{WIMP}}$$

# Other Mechanisms

- Gravitinos are the original SUSY dark matter

Pagels, Primack (1982)  
Weinberg (1982)  
Krauss (1983)  
Nanopoulos, Olive, Srednicki (1983)

Khlopov, Linde (1984)  
Moroi, Murayama, Yamaguchi (1993)  
Bolz, Buchmuller, Plumacher (1998)  
...

Old ideas:

- Gravitinos have thermal relic density

$$\Omega_{\tilde{G}} < 1 \Rightarrow m_{\tilde{G}} < 1 \text{ keV}$$

- For DM, require a new, fine-tuned energy scale

- Weak scale gravitinos diluted by inflation, regenerated in reheating

$$\Omega_{\tilde{G}} < 1 \rightarrow T_{\text{RH}} < 10^{10} \text{ GeV}$$

- For DM, require a new, fine-tuned energy scale



# SuperWIMP Signals

Typical reactions:

- A) Signals too strong; this scenario is completely excluded
- B) Signals too weak; this scenario is possible, but completely untestable

Can't both be right – in fact both are wrong!

# SuperWIMP Signals

- SuperWIMPs escape all conventional DM searches
- But late decays  $\tilde{\tau} \rightarrow \tau \tilde{G}$ ,  $\tilde{B} \rightarrow \gamma \tilde{G}$ , ..., have cosmological consequences
- Assuming  $\Omega_{\tilde{G}} = \Omega_{\text{DM}}$ , signals determined by 2 parameters:

$$m_{\tilde{G}}, m_{\text{NLSP}}$$

Lifetime

$$\Gamma(\tilde{\ell} \rightarrow \ell \tilde{G}) = \frac{1}{48\pi M_*^2} \frac{m_{\tilde{\ell}}^5}{m_{\tilde{G}}^2} \left[1 - \frac{m_{\tilde{G}}^2}{m_{\tilde{\ell}}^2}\right]^4$$

$$\Gamma(\tilde{B} \rightarrow \gamma \tilde{G}) = \frac{\cos^2 \theta_W}{48\pi M_*^2} \frac{m_{\tilde{B}}^5}{m_{\tilde{G}}^2} \left[1 - \frac{m_{\tilde{G}}^2}{m_{\tilde{B}}^2}\right]^3 \left[1 + 3 \frac{m_{\tilde{G}}^2}{m_{\tilde{B}}^2}\right]$$

Energy release


$$\zeta_i = \varepsilon_i B_i Y_{\text{NLSP}}$$

i = EM, had

$$Y_{\text{NLSP}} = n_{\text{NLSP}} / n_{\gamma}^{\text{BG}}$$

# Big Bang Nucleosynthesis

Late decays may modify light element abundances



Fields, Sarkar, PDG (2002)

After WMAP

- $\eta_D = \eta_{\text{CMB}}$
- Independent  ${}^7\text{Li}$  measurements are all low by factor of 3:

$${}^7\text{Li}/\text{H} = 1.5_{-0.5}^{+0.9} \times 10^{-10} \quad (95\% \text{ CL}) \quad [27]$$

$${}^7\text{Li}/\text{H} = 1.72_{-0.22}^{+0.28} \times 10^{-10} \quad (1\sigma + \text{sys}) \quad [28]$$


$${}^7\text{Li}/\text{H} = 1.23_{-0.32}^{+0.68} \times 10^{-10} \quad (\text{stat} + \text{sys}, 95\% \text{ CL}) \quad [29]$$

- ${}^7\text{Li}$  is now a serious problem

Jedamzik (2004)

# BBN EM Constraints


- NLSP = WIMP  $\rightarrow$  Energy release is dominantly EM (even mesons decay first)
- EM energy quickly thermalized, so BBN constrains  $(\tau, \zeta_{EM})$
- BBN constraints weak for early decays: hard  $\gamma$ ,  $e^-$  thermalized in hot universe
- Best fit reduces  ${}^7\text{Li}$ : 😊



Cyburt, Ellis, Fields, Olive (2002)

# BBN EM Predictions

- Consider  $\tilde{\tau} \rightarrow \tilde{G} \tau$  (others similar)
- Grid: Predictions for  
 $m_{\tilde{G}} = 100 \text{ GeV} - 3 \text{ TeV}$  (top to bottom)  
 $\Delta m = 600 \text{ GeV} - 100 \text{ GeV}$  (left to right)
- Some parameter space excluded, but much survives
- SuperWIMP DM naturally explains  ${}^7\text{Li}$  !



Feng, Rajaraman, Takayama (2003)

# BBN Hadronic Constraints

- BBN constraints on hadronic energy release are severe.

Jedamzik (2004)

Kawasaki, Kohri, Moroi (2004)

- For neutralinos, hadrons from

$$\chi \rightarrow Z\tilde{G}, h\tilde{G}$$



exclude  $\chi = \tilde{W}, \tilde{h}, \tilde{B}$ . Only  $\chi = \tilde{\gamma}$  and  $\Delta m < m_Z$  are ok.

- For sleptons, cannot neglect subleading decays:

$$\tilde{l} \rightarrow lZ\tilde{G}, \nu W\tilde{G}$$

$$\tilde{\nu} \rightarrow \nu Z\tilde{G}, lW\tilde{G}$$

# BBN Hadronic Predictions



Feng, Su, Takayama (2004)

Despite  $B_{\text{had}} \sim 10^{-5} - 10^{-3}$ , hadronic constraints are leading for  $\tau \sim 10^5 - 10^6$ , must be included

# Cosmic Microwave Background

- Late decays may also distort the CMB spectrum

- For  $10^5 \text{ s} < \tau < 10^7 \text{ s}$ , get “ $\mu$  distortions”:


$$\frac{1}{e^{E/(kT)+\mu} - 1}$$

$\mu=0$ : Planckian spectrum

$\mu \neq 0$ : Bose-Einstein spectrum

Hu, Silk (1993)



- Current bound:  $|\mu| < 9 \times 10^{-5}$
- Future (DIMES):  $|\mu| \sim 2 \times 10^{-6}$



Feng, Rajaraman, Takayama (2003)



# SUSY Spectrum ( $\Omega_{\tilde{G}} = \Omega_{\text{DM}}$ )



Feng, Su, Takayama (2004)

$\Omega_{\tilde{G}} = (m_{\tilde{G}} / m_{\text{NLSP}}) \Omega_{\text{NLSP}}$  results – see Su’s talk

# Model Implications

- We've been missing half of parameter space.  
For example, mSUGRA should have **6** parameters:



$$\{ m_0, M_{1/2}, A_0, \tan\beta, \text{sgn}(\mu), m_{3/2} \}$$

$\tilde{G}$  not LSP

$\Omega_{\text{LSP}} > 0.23$  excluded

$\tilde{G}$  LSP

$\Omega_{\text{NLSP}} > 0.23$  ok



# Collider Phenomenology

Drees, Tata (1990)

Goity, Kossler, Sher (1993)

Feng, Moroi (1996)

Hoffman, Stuart et al. (1997)

Acosta (2002)

...

- Each SUSY event produces 2 metastable sleptons  
Spectacular signature: highly-ionizing charged tracks


Current bound (LEP):  $m_{\tilde{\tau}} > 99 \text{ GeV}$

Tevatron Run II reach:  $m_{\tilde{\tau}} \sim 150 \text{ GeV}$

LHC reach:  $m_{\tilde{\tau}} \sim 700 \text{ GeV}$  in 1 year

# Slepton Trapping

- Sleptons can be trapped then moved to a quiet environment to observe decays
- LHC:  $10^6$  sleptons/yr possible. Slow sleptons are isotropic. By optimizing trap location and shape, can catch  $\sim 100$ /yr in  $100 \text{ m}^3$  we
- LC: tune beam energy to produce slow sleptons



Smith et al., in preparation

# Measuring $m_{\tilde{G}}$ and $M_*$

- Recall:

$$\Gamma(\tilde{\ell} \rightarrow \ell \tilde{G}) = \frac{1}{48\pi M_*^2} \frac{m_{\tilde{\ell}}^5}{m_{\tilde{G}}^2} \left[ 1 - \frac{m_{\tilde{G}}^2}{m_{\tilde{\ell}}^2} \right]^4$$

- Measurement of  $\Gamma \rightarrow m_{\tilde{G}}$ 
  - $\rightarrow \Omega_{\tilde{G}}$ . SuperWIMP contribution to dark matter
  - $\rightarrow F$ . Supersymmetry breaking scale
  - $\rightarrow$  BBN in the lab
- Measurement of  $\Gamma$  and  $E_l \rightarrow m_{\tilde{G}}$  and  $M_*$ 
  - $\rightarrow$  Precise test of supergravity: gravitino is graviton partner
  - $\rightarrow$  Measurement of  $G_{\text{Newton}}$  on fundamental particle scale
  - $\rightarrow$  Probes gravitational interaction in particle experiment

# Recent Related Work

- SuperWIMPs in universal extra dimensions  
Feng, Rajaraman, Takayama, hep-ph/0307375
- Motivations from leptogenesis  
Fujii, Ibe, Yanagida, hep-ph/0310142
- Impact on structure formation  
Sigurdson, Kamionkowski, astro-ph/0311486
- Analysis in mSUGRA  
Ellis, Olive, Santoso, Spanos, hep-ph/0312062  
Wang, Yang, hep-ph/0405186
- Collider gravitino studies  
Buchmuller, Hamaguchi, Ratz, Yanagida, hep-ph/0402179, hep-ph/0403203

# Summary

|                                  | WIMPs | superWIMPs                    |
|----------------------------------|-------|-------------------------------|
| Well-motivated stable particle?  | Yes   | Yes                           |
| Naturally correct relic density? | Yes   | Yes                           |
| Detection promising?             | Yes   | Yes<br>${}^7\text{Li}$ signal |

SuperWIMPs – a new class of particle dark matter with completely novel implications